

VALIDITY OF INTRACEREBRAL HAEMORRHAGE SCORE IN PREDICTING PROGNOSIS OF HEMORRHAGIC STROKE

Dr. Muhammad Aslam Rind¹, Dr. Muzaffar sheikh², Dr. Manzoor Ali lakhair³, Dr. Maria Nazir⁴, Dr. Sadia Sheikh⁵

¹. Senior Registrar Department of Medicine

². Professor Department of medicine

³. Associate professor Department of Neurology

⁴. Medical officer

Correspondence to: Dr. Manzoor Ali lakhair Associate professor Department of neurology Liqueat university of medical and health sciences LUMHS Email: manzoor.ali@lumhs.edu.pk
Date of Submission: August 22, 2015 **Date of Revision:** October 09, 2015, **Date of Acceptance:** October 25, 2015

ABSTRACT

Background: Stroke is a common neurological disorder, leading cause of death and disability worldwide. Intra cerebral hemorrhage comprises of 10-15% of all strokes but associated with high risk of morbidity and mortality than cerebral infraction. The ICH Score is a good prognostic tool calculated very easily, when it is used to predict 30 day mortality. **Objective:** To determine the validity of Intracerebral hemorrhagic score in predicting prognosis (adverse outcome or death) in patients of hemorrhagic stroke. **Material & Methods:** This case series study was carried out at department of medicine Liaquat medical University Hospital Jamshoro and Hyderabad from June 16th 2012 to December 15th 2012. Eighty nine 89 patients were enrolled in the study, non-probability consecutive sampling technique was applied and data was collected after inclusion and exclusion criteria and structured proforma were designed for study. **Results:** Out of 89 patients 56 (62.9%) were male and mean age of the patients in our study was 65.14 years. In 32(36%) patients ICH score was ≥ 3 while 57 (64%) patients had ICH score was < 3 . Out of total 32 patients who had ICH score ≥ 3 , Adverse outcome i.e. death occurred in 25 (78.12%) patients and males were 16(80%) and 9(75%) were female patients. **Conclusion:** Frequency of adverse outcome (death) was found in 28.1% while ICH score predicted adverse outcome (death) in 36.1%. ICH score correctly predicted 77.77% adverse outcome, therefore we can conclude that ICH score is useful scale to predict mortality in patients suffering from intracerebral hemorrhage.

KEY WORDS Intracerebral Hemorrhage, Intra cerebral hemorrhage score, Glasgow Coma Scale, Mortality

INTRODUCTION

Stroke is defined as the sudden onset of focal neurological deficit due to a vascular cause. Stroke results from lack of blood flow to an area of the brain. Stroke is classified into ischemic and hemorrhagic stroke. Hemorrhagic stroke may be either cerebral haemorrhage or intra cranial haemorrhage and cerebral haemorrhage is either intra parenchymal haemorrhage or intra ventricular haemorrhage. Intracranial haemorrhage classified in to extra Dural hematoma, subdural hematoma and subarachnoid haemorrhage. Stroke is the common neurological disorder, leading cause of death in developing as well as in developed countries, prevalence in us is about 7 million (3.0%) and incidence varies according to age, sex, ethnicity and socioeconomic status [1]. Worldwide 15 million people suffer from stroke each year, one third die and one-third are left permanently disabled[1][2]. Stroke sub types vary in different parts of the world between different races but increased proportion of intracerebral haemorrhage and lacunar infarcts have been reported in Asia [3][4]. Intra cerebral haemorrhage comprises of 10-15% of all strokes but has a

much higher risk of morbidity and mortality than cerebral infraction or subarachnoid haemorrhage (SAH) [5]. Despite advances in the treatment of cerebral infraction and SAH, all hematomas are not surgically accessible or their drainage doesn't always improve outcome [6] and no known medical management strategy (like tissue plasminogen activator in ischemic stroke) in case of Intracerebral haemorrhage which improves the outcome of ICH so there should be some adequate scoring system which will help to predict prognosis of ICH [7]. Standardization is a major issue for triage and subsequent management of such patients [8-9]. Studies on ICH treatment has used a variety of selection criteria for patient's inclusion but inconsistency of selection criteria across studies emphasize that there is no standard or widely accepted early prognostic model or clinical grading scale for ICH that analogous to those used for cerebral infraction, SAH, or traumatic brain injury.[10] [11][12]. The expected frequency of mortality in intracerebral haemorrhage is around 45% [13]. The prognosis of hemorrhagic stroke varies depending on the site and size of the haemorrhage. Lower Glasgow Coma Scale scores are associated with poorer

prognosis and higher mortality rate. A larger volume of blood at presentation is associated with a poorer prognosis. Growth of the hematoma volume is associated with a poorer functional outcome and increased mortality rate. Non aneurysmal perimesencephalic has a less severe clinical course and in general better prognosis. The ICH Score utilizes combination of clinical information and CT scan recording [5], can be calculated very easily and this may be a good prognostic tool when used to predict 30 day mortality in patients with intracerebral haemorrhage. In the prediction of mortality and short-term outcomes in patients with intracerebral haemorrhage (ICH), the ICH score is a simple, reliable, broadly accepted grading scale. The score is a composite of admission level of consciousness (Glasgow Coma Scale score), age, and initial neuroimaging findings (hematoma volume, location, and intra ventricular haemorrhage). Objective of this study to determine the frequency of adverse outcome (death) correctly predicted by the ICH scores in patients suffering from intracerebral haemorrhage. The ICH score was developed retrospectively and until now had not been validated for prediction of long-term functional outcome

MATERIAL & METHOD

This case series study was carried out at department of medicine Liaquat medical University Hospital Jamshoro and Hyderabad for a period of six months from June 16, 2012 to December 15, 2012. All the admitted patients with stroke, suspected of having ICH were considered for enrollment in the study after obtaining the informed consent and according to the predetermined inclusion and exclusion criteria. Eighty nine 89 patients were enrolled in this case series study and non-probability consecutive sampling technique was applied, Structured proforma was designed to fill the data on the basis of inclusion and exclusion criteria. Following is the inclusion criteria, Patients of stroke in adult age and confirmed cases of intra cerebral haemorrhage (Intra cerebral haemorrhage is defined as the presence of clinical focal neurological deficit >24 hours, altered sensorium, fluctuating consciousness and confirmed by CT scan findings by visualization of haemorrhage with in the cranial vault as per ICD 10, 161.x) were included in study. Patients with following criteria were excluded, Age <14 was excluded Patients of stroke other than intra cerebral haemorrhage, with meningo-encephalitis and intra cranial space occupying lesion were also excluded from study after clinical and radiological evaluation. Data was analyzed on SPSS version 20. Mean and standard deviation was calculated for age, RBS, ICH score, Mean arterial pressure and systolic blood pressure. Frequencies were calculated for categorical variables like gender, GCS score category, ICH volume category, IVH, Infratentorial origin of ICH, age category and mortality. Stratification was done about age

and gender Other clinical and radiological parameters of enrolled patients in this study were also noted like Initial blood pressure, pulse pressure, mean arterial pressure and Glasgow Coma Scale (GCS). On CT scan Brain intra cerebral haemorrhage volume (ICH hematoma volume was measured on the initial head CT scan with the use of the ABC/2 method, in which A is the greatest diameter on the largest haemorrhage slice, B is the diameter perpendicular to A and C is the approximate number of axial slices with haemorrhage multiplied by the slice thickness) was recorded and presence or absence of IVH was also noted as a component of ICH score. Patients were followed up to 30 days after hospital admission. Primary outcome was taken as death within 30 days of admission.

RESULTS

Mean age of the patients in our study was 65.14 years with the standard deviation of ± 6.43 years. The minimum age was 55 years, while the maximum age was 81 years, as shown in table 1.

Table 1: Age of the patient n=89

Age of the patients (Years)	Mean	S.D	Minimum	Maximum
	65.14	± 6.43	55	81

Gender distribution shows that out of total 89 patients 56 (62.9%) were male while 33 (37.1%) were female patients as shown in figure 1.




Figure 1: Gender Distribution

Regarding age distribution 83 (93.3%) patients were <80 years of age. (Figure 2).




Figure 2: Age distribution

Mean RBS, Intracerebral Haemorrhage score, mean arterial pressure and systolic blood pressure were found to be 145.88 ± 60.38 mg/dl, 4.49 ± 0.740 , 135.64 ± 27.36 mm/Hg, and 172.13 ± 22.18 mm/Hg as shown in tables 2, 3, 4 and 5 respectively.

RBS (mg/dl)	Mean	S.D	Minimum	Maximum
	145.88	60.38	85	300

Table 2: RBS n=89

ICH Score	Mean	S.D	Minimum	Maximum
	4.49	0.74	3	5

Table 3: ICH Score n=89

MAP (mm/hg)	Mean	S.D	Minimum	Maximum
	135.64	± 27.36	92	170

Table 4: Mean Arterial Pressure n=89

SBP	Mean	S.D	Minimum	Maximum
	172.13	± 22.18	140	220

Table 5: Systolic Blood Pressure n=89

In 71(79.8) patients Glasgow coma scale (GCS) was between 3-4, while in 18 (20.2%) patient's GCS was between 5-12, as shown in figure 3.




Figure 3: Glasgow coma scale

In this study Intracerebral haemorrhage volume was ≥ 30 cm³ in 46(51.7%) patients as shown in figure 4 and 57(64%) patients had intra ventricular haemorrhage (IVH), while only 10 (11.2%) patients had infra tentorial haemorrhage, as shown in Figure 5 and 6.




Figure 4: ICH Volume




Figure 5: IVH IVH=Intra ventricular haemorr




Figure 6: Infra tentorial haemorrhage

In our study 32(36%) patients had ICH score ≥ 3 while 57 (64%) patients had ICH score < 3 as shown in figure 7.




Figure 7: ICH Score ICH= Intracerebral haemorrhage

Out of total 32 patients who had ICH score ≥ 3 , actual Adverse outcome, i.e. death occurred in 25 (78.12%) patients whereas 7(21.85%) patients remained alive as mentioned in Figure 8.




Figure 8: Mortality in patients with ICH score > 3

Out of 17 patients with age < 80 years mortality occurred in 11 (64.70%) patients remaining 6(35.29%) were alive, whereas 14 (93.33%) patients had a mortality with age ≥ 80 years, as shown in Table 6

Age Distribution	Mortality		Total
	Yes	No	
< 80 years	11(64.70)	6(35.29)	17(100)
≥ 80 years	14(93.33)	1(6.66)	15(100)
Total	25(78.12)	7(21.87)	32(100%)

Table 6: Age categories and mortality in patients with ICH score ≥ 3 n=

Regarding gender distribution mortality with ICH score

≥ 3 , male patients were 16(80%) and 9(75%) patients were females as shown in Table 7.

Gender Distribution	Mortality		Total
	Yes	No	
Male	16 (80)	4 (20)	20 (100)
Female	9 (75)	3 (25)	12 (100)
Total	25(78.12)	7(21.87)	32(100%)

Table 7: Gender Distribution and mortality in patients with ICH score ≥ 3 n=32

DISCUSSION

Intracerebral haemorrhage (ICH) causing 10 to 15% of first-ever stroke, with 30-day mortality rate of 35 to 52% and half of the deaths reported within the first 2 days.[14] [15][16]. Mean age of the patients in our study was 65.14 ± 6.43 years. Similar results were published by patriota et al in 2009 , mean age in their study was 67.7 ± 11.2 years [17]. Mean age of the patients in the study of Weimar et al was 67.4 ± 12 years [18]. In our study 37.1% patients were females while most of the patients were males i.e. 62.9% but 51.4% patients were females in the study of Patriota et al [17]. Gender distribution in the study of Weimar et al shows 43.9% patients were females in their study [18]. Mean RBS and systolic blood pressure in our study was found to be 145.88 ± 60.38 mg/ and 172.13 ± 22.18 mm/Hg respectively and almost Similar results were also reported 144.1 ± 62.8 mg/dl and 173.8 ± 35.7 mm/Hg by Patriota et al[17]. Out of total 32 patients who had ICH score was ≥ 3 actual adverse outcome i.e. death occurred in 25 (78.12%) patients, whereas in contrast the mortality was found to be 37.8% in another study [17]. In few studies ICH score was previously been validated in an independent dataset for prediction of good outcome but mortality rate during the initial hospitalization was high (95 of 243 patients) and higher ICH score was associated with a lower likelihood of favorable outcome [18][19]. The expected frequency of mortality in Intracerebral haemorrhage is around 45%[13], Hemphill et al conducted a landmark study in which ICH was validated[5] and 161 patients with ICH score 5 were died whereas thirty day mortality rate for patients with ICH scores of 1,2,3 and 4 were 13%,26%, 72% and 97% respectively[.5]

CONCLUSION

In this study frequency of adverse outcome (death) was found in 28.1% while ICH score predicted adverse outcome (death) in 36.1%. ICH score correctly predicted 77.77% adverse outcome, therefore we can conclude that the ICH score is a simple, reliable, broadly accepted grading scale. This study confirms the literature data about high mortality in ICH and reinforces the external validation of ICH Score as good prognostic tool to predict the adverse outcome in intra cerebral hemorrhagic stroke.

REFERENCE

1. Roger et al. AHA Heart Disease and Stroke Statistics 2011 update: a report from the American Heart Association. *Circulation* 2011; 123:e18-e209.
2. The Atlas of heart disease and stroke, WHO 2004. http://www.who.int/cardiovascular_diseases/en/cvd_atlas_15_burden_stroke.pdf (Mackay J, Mensah G: The Atlas of Heart Disease and Stroke. Geneva, Switzerland, World Health Organization, 2004)
3. Avorn J, Kesselheim A. A haemorrhage of off-label use. *Ann Intern Med.* Apr 19 2011; 154(8) :566-7.
4. Yank V, Tuohy CV, Logan A C, et al. Systematic Review: Benefits and Harms of In-Hospital Use of Recombinant Factor VIIa for Off-Label Indications. *Ann InternMed.* Apr 19 2011; 154(8):529-40.
5. Broderick J, Connolly S, Feldmann E, Hanley D, Kase C, and Krieger D, et al. Guidelines for the management of spontaneous intracerebral haemorrhage in adults: 2007 update: a guideline from the American Heart Association/American Stroke Association Stroke Council, High Blood Pressure Research Council, and the Quality of Care and Outcomes in Research Interdisciplinary Working Group. *Circulation.* Oct 16 2007; 116(16) :e391-413.
6. Shiber JR, Fontane E, Adewale A. Stroke registry: hemorrhagic vs ischemic strokes. *Am J Emerg Med.* Mar 2010; 28(3):331-3.
7. Lloyd-Jones D, Adams R, Carnethon M, De Simone G, Ferguson TB, Flegal K, et al. Heart disease and stroke statistics--2009 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation.* Jan 27 2009; 119(3):480-6.
8. Adams HP Jr, Bendixen BH, Kappelle LJ, BillerJ, Love BB, Gordon DL, et al. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in Acute Stroke Treatment. *Stroke.* Jan 1993; 24(1):35-41.
9. Thrift AG, Dewey HM, MacDonnell RA, McNeil JJ, and Donnan GA. Incidence of the major stroke subtypes: initial findings from the North East Melbourne stroke incidence study (NEMESIS). *Stroke.* Aug 2001; 32(8):1732-8.
10. Donnan GA, Fisher M, Macleod M, Davis SM. Stroke. *Lancet.* May 10 2008; 371(9624) :1612-23.
11. Mullins ME, Lev MH, Schellingerhout D, Gonzalez RG, Schaefer PW. Intracranial haemorrhage complicating acute stroke: how common is hemorrhagic stroke on initial head CT scan and how often is initial clinical diagnosis of acute stroke eventually confirmed?. *AJNR Am J Neuroradiology.* Oct 2005; 26(9):2207-12.
12. Nighoghossian N, Hermier M, Adeleine P, Blanc-Lasserre K, Derex L, Honnorat J, et al. Old micro bleeds are a potential risk factor for cerebral bleeding after ischemic stroke: a gradient-echo T2*-weighted brain MRI study. *Stroke.* Mar 2002; 33(3):735-42.
13. Kim EY, Na DG, Kim SS, Lee KH, Ryou JW, Kim HK. Prediction of hemorrhagic transformation in acute ischemic stroke: role of diffusion-weighted imaging and early parenchymal enhancement. *AJNR Am J neuroradiology.* May 2005; 26(5):1050-5.
14. Broderick JP, Brott T, Tom sick T, et al. Intracerebral haemorrhage more than twice as common as subarachnoid haemorrhage. *J Neurosurg* 1993; 78:188-191.
15. Anderson CS, Chakera TM, Stewart-Wynne EG, et al. Spectrum of primary intracerebral haemorrhage in Perth, Western Australia, 1989-90: incidence and outcome. *J Neurol Neurosurg Psychiatry* 1994; 57:936-940.
16. Counsell C, Boonyakarnkul S, Dennis M, et al. Primary Intracerebral haemorrhage in the Oxfordshire Community Stroke Project, 2: prognosis. *Cerebrovascular Dis* 1995; 5:26-34.
17. Patriota GC, da Silva JM, Barcellos AC, de Sousa JB, Toledo DO, Pinto F et al. Determining ICH Score: can we go beyond? *Arq Neuropsiquiatr* 2009; 67(3-A):605-608.
18. Weimar C, Benemann J, Diener HC. Development and validation of the Essen Intracerebral Haemorrhage Score. *J Neurol*

- Neurosurg Psychiatry 2006; 77:601–605.
19. Hemphill JC III et al. Prospective validation of the ICH score for 12-month functional outcome. Neurology 2009 Oct 6; 73:1088.

Conflict of interest: Author declares no conflict of interest.

Funding disclosure: Nil

Author's contribution:

Dr. Muhammad Aslam Rind: Protocol Writing Study Concept & Design, Data Collection, Data analysis, manuscript writing & Manuscript Review.

Dr. Muzaffar sheikh: Protocol Writing Study Concept & Design, Data Collection, Data analysis, manuscript writing & Manuscript Review.

Dr. Manzoor Ali lakhair: Protocol Writing Study Concept & Design, Data Collection, Data analysis, manuscript writing & Manuscript Review.

Dr. Maria Nazir: Data Collection, Data analysis, manuscript writing & Manuscript Review.

Dr. Sadia Sheikh: Data Collection, Data analysis, manuscript writing & Manuscript Review.